

Protein Feeding Strategies for Dairy Replacement Heifers

By Patrick C. Hoffman
Dairy Specialist
Department of Dairy Science
University of Wisconsin-Madison
Marshfield Agricultural Research Station
8396 Yellowstone Drive
Marshfield, WI 54449
715-387-2523
fax 715-387-1723
pchoffma@facstaff.wisc.edu



Protein Feeding Strategies For Dairy Replacement Heifers

Before beginning a discussion of protein feeding strategies for dairy replacement heifers, a discussion of the fundamentals of replacement heifer growth is required. It is relevant because a basic understanding of growth fundamentals provides the foundation for understanding protein requirements of dairy replacement heifers. Most producers, consultants, veterinarians, educators, etc., are familiar with heifer growth charts which define optimum weight, height to age relationships.

Growth charts are invaluable when monitoring the effectiveness of a replacement heifer management program. While invaluable, a heifer growth chart does not, however, provide insight into the complexities of replacement heifer growth.

Often producers, consultants, etc., desire to produce heifers differently than those defined on a growth chart. Reducing the calving age via accelerated growth is a common objective. Accelerating growth rate has a desired effect in increasing the rate of body protein deposition. Heifers fed diets with enough energy to support 2.2 lbs/d of average daily gain (ADG) will have accelerated body protein deposition as compared to heifers fed diets to gain 1.75 lbs/d.

The differences in body protein deposition between heifers gaining 2.2 vs 1.75 lbs/d are more pronounced in the first year of life. Logically, if accelerated growth enhances body protein deposition, increases in dietary protein supply would be required. There is, however, a major biological obstacle to accelerated growth of replacement heifers. Accelerated growth does increase the rate of body protein deposition, but it also increases the rate of body fat deposition.

Replacement heifers gaining 2.2 lbs/d deposit more fat than heifers gaining 1.75 lbs/d. Differences in body fat deposition become more pronounced in the second year of life. Thus in a way the positive aspects of increased body protein deposition in the first year of life via accelerated

growth are negated by negative effects of fat deposition in the second year of life. These combined factors result in the production of replacement heifers which may weigh the same (e.g., 1400 lbs at calving), calve at different ages (20 vs 24 months), but have different body compositions.

Replacement heifers reared to 1400 lbs by gaining 1.75 lbs/d would theoretically have 22.9% body fat vs 36.9% for heifers reared at 2.2 lbs/d. Conversely, 1400 lb heifers which gained 1.75 lbs/d would have 17.4% body protein as compared to 14.4% for heifers gaining 2.2 lbs/d. Differences in body composition at calving have been demonstrated (10) to result in critical differences in first lactation milk yield.

Over the past decade, researchers have been investigating protein feeding strategies that may override some of these fundamentals of replacement heifer growth, allowing the production of replacement heifers at an earlier age without compromising body composition issues.

Crude Protein vs. Heifer Growth

Calf Starters – High rates of true body protein deposition occurs at very young ages (figure 2) and necessitates that pre-post weaning calf starters be relatively high in crude protein (CP). Minnesota workers (2) fed Holstein calves grain starters containing 15.0, 16.8, 19.6, or 22.4% CP. Overall body weight gains from day 4 to 56 averaged 1.20, 1.23, 1.37, and 1.36 lbs/d, respectively. Throughout the experiment, starter consumption tended to increase and CP content of the starter increased.

The researchers concluded that 19.6 % CP in starters was adequate. Wisconsin workers (11) fed Holstein calves starters containing 20.0 or 25.0% CP. Starters were first offered on day 21 prior to weaning at day 26. Calf growth rates were nearly identical (.60 lbs/day) for calves fed starters containing 20.0 or 25.0% CP. Likewise, the researchers concluded that 20% CP in the starter was adequate.



These data (2,11) suggest that starters containing 19 to 20% CP on a dry matter basis appear to be adequate in supporting calf growth rates. The reader is cautioned that nutrient compositions of commercial calf starters are listed on an as fed basis. Therefore commercial calf starters listing 17 to 18% CP on the feed tag are actually 19 to 20% on the dry matter basis.

Heifer Diets

Surprisingly, there are few recent studies that have evaluated CP feeding regimens for older heifers. Canadian workers (4) fed heifers from 71 to 126 days 13.3, 16.6, or 19.2% CP followed by 10.9, 14.0, and 17.3% CP from 127 to 182 days. Results suggest that weight and height growth of Holstein heifers was maximized when diets contained 16.6% CP (71 to 126 days) followed by 14% CP (127 to 182 days). Italian workers (14) fed Holstein heifers weighing 220 to 440 lbs diets containing 16.7 or 13.7% CP. Heifers fed 16.7% CP gained 1.72 lbs/d as compared to 1.64 lbs/d for heifers fed 13.7% CP.

In a second phase of the study, heifers were fed either 12.5% or 10.4% CP from 440 lbs to 660 lbs. Heifers fed 12.5% CP gained 1.66 lbs/d as compared to 1.54 lbs/d for heifers fed 10.4% CP. Beltsville workers (23) fed Holstein heifers from 400 to 736 lbs alfalfa based diets containing 22.0% CP vs corn silage based diets containing 15.6% CP. Gains were similar between the diets at 1.91 and 1.98 lbs/d. In addition, there were no appreciable differences in body protein gain. In a recent study (9), we fed Holstein heifers weighing 880 lbs diets

containing 8.1, 10.7, 12.7, and 15.0% CP, respectively (Table 1).

There were no significant effects on ADG, but wither height, hip width, and heart girth increased linearly with increasing dietary protein, but these were slight decreases or no change in these measurements when heifers were fed 15.0% CP. In addition, blood protein parameters and nitrogen balance of heifers were measured. Blood urea nitrogen levels were extremely low when heifers were fed diets containing 8.1 or 10.7% CP. Nitrogen (CP) retention was maximized in heifers fed diets containing 13.7% CP.

While data are limited, these studies (4, 9, 14, 23) suggest Holstein heifers from 200 to 500 lbs be fed diets containing approximately 15 to 17% CP, heifers 500 to 800 lbs, 14 to 15% CP, and heifers >800 lbs, 13% CP. These guidelines are similar to, but slightly higher than, current NRC (13) recommendations.

Bypass Protein and Heifer Growth

Over the past decade there have been a number of studies conducted to evaluate the potential of feeding bypass protein to augment heifer growth and body protein deposition. The following discussion focuses on those efforts.

Calf Starters – Pennsylvania workers (18) fed calves from birth to 25 weeks starters containing 33, 37, and 46% bypass protein (% of CP). There were no appreciable differences in calf growth rates between the treatments. Slaughter data also suggested no body protein deposition advantages to feeding starters higher in bypass protein because carcass fat and protein levels were nearly identical between the treatments.

In contrast, New Hampshire (17) did observe increased growth rates (1.64 vs 1.54 lbs/d) when starters containing 46% bypass protein were compared to starters containing 36% bypass protein. Kansas State workers (1) observed benefits in calf growth when soybeans or corn were roasted, thereby increasing the bypass protein content. Based on these data (1, 17, 18), it is very difficult to ascertain what level of bypass protein may be beneficial to calves. This issue is probably

Table 1. Growth, blood metabolites, and nitrogen balance of 880-lb. Holstein heifers fed 8.1%, 10.7%, 12.7%, or 15.0% CP diets.

	(Diet CP)			
	8.1%	10.7%	12.7%	15.0%
Growth				
ADG, lbs/d	1.40	1.52	1.63	1.54
Height, in/120 d	1.50	2.68	3.54	2.83
Hip width, in/120 d	1.85	2.16	2.40	2.32
Heart girth, in/120d	4.41	5.55	6.41	6.41
Blood Metabolites				
BUN mg/dl	0.4	5.6	9.6	12.4
N-Balance				
N-retention g/d	12.0	17.0	23.0	22.0

more related to the quality of amino acid availability of the bypass protein source. For example, Pennsylvania workers (18) used blood meal to increase starter bypass contents while New Hampshire (17) and Kansas State (1) workers used heat treated soy based proteins. Differences in bypass source could account for the differences in heifer performance between experiments.

While more information is needed, there may be some advantages to feeding calves starters that contain some additional bypass. While speculative, heat treated plant proteins may be more effective than animal proteins in calf starters.

Heifer Diets

There have been a number of experiments conducted evaluating the effects of increased dietary bypass protein on heifer growth and performance. Describing the results of each trial would be tedious, so the author has constructed a summary table for ease of interpretation (Table 2).

In general, supplementing bypass protein to dairy replacement heifers has not resulted in signif-

icant or consistent improvement in heifer growth or feed efficiency. Results from two of the trials (3, 6) may be confounded by interactions with energy which also were part of the experimental protocol. This leaves one trial (20) where a true growth response was observed to supplementary bypass protein. The positive responses observed in this

Table 3. Results of feeding high CP, high energy diets to Holstein heifers.

<u>item</u>	<u>control</u>	<u>high energy/ protein</u>
Slaughter Study		
Slaughter BW	740	873
ADG, lbs/day	1.7	2.62
Carcass Protein, %	17.4	16.3
Carcass Fat, %	16.6	24.8
Mammary Tissue, g	401	408
Lactation Study		
Calving Age, mos.	23.6	20.7
Milk, lbs/day	64.2	56.4

Table 2. A review of recent literature regarding bypass protein supplementation.

<u>refer- ence</u>	<u>weight</u>	<u>bypass source</u>	<u>1</u>	<u>diet bypass level</u>			<u>5</u>	<u>feed growth resp.</u>	<u>efficiency response</u>
				<u>2</u>	<u>3</u>	<u>4</u>			
21	200-600	Animal Proteins	30	40				No	No
19	550-1,100	Animal Proteins	34	42				No	N/A
6	330	Extruded Soybeans	34.5	35.4				No	N/A
6	330	Extruded Soybeans	28.5	29.1				Yes	N/A
3	220	Corn Gluten Meal	36	53				Yes	Yes
5	400-900	Animal Proteins	30	50				No	Yes
20	500	Animal Proteins	31	43	50	55		Yes	Yes
12	400-1,000	Animal Proteins	28	33	30	33	34	No	No
		Animal Proteins	31	38	35	39	40	No	No



study are somewhat inconclusive, however, as only weight gains were improved. Growth of wither height, hip height, and heart girth were not improved by increasing bypass protein content of replacement heifer diets. The body of recent research work suggests major alteration of bypass protein in replacement heifer diets is probably not warranted.

Dietary Protein to Energy Ratio

Recently, Michigan researchers (22) have proposed that dietary protein to energy ratios are an important factor in replacement heifer diets. This is a logical hypothesis because as dietary energy is increased to accelerate growth, body protein deposition rate is increased, thereby increased dietary protein would be required. This concept was used to evaluate the often conflicting and confusing studies on mammary development. Michigan workers (22) evaluated 11 mammary development experiments and observed that mammary development was inversely related ($r^2 = 0.71$) with the CP to energy (ME) ratio of the diet.

Further, the CP:ME ratio of the diet accounted for 76% of the variation in milk yield from these studies. Simplistically, these data suggest that mammary development and milk yield

are impaired when heifers are fed low protein, high energy diets. These data also suggest when high energy diets are fed, increasing dietary protein may be warranted. In recent studies (15, 16), Michigan workers tested this hypothesis. Heifers weighing 275 lbs were fed a diet containing .53 Mcals/lb of NEM and 16.3% CP or a diet containing .83 Mcals of NEM and 19.3% CP. Summary results of this study are presented in Table 3.

The encouraging part of this study was that body

Table 4. Ration specifications for Holstein replacement heifers.^{1,2}

Item	Unit	(heifer weight, lbs)							
		175	375	575	775	975	1175	1375	
Intake	lbs/d	6.3	11.3	15.3	18.3	22.4	26.4	30.4	
Energy									
TDN ³	% of DM	76.0	69.0	66.0	64.0	64.0	64.0	68.0	
ME	Mcal/lb	1.30	1.18	1.12	1.12	1.08	1.08	1.20	
Protein									
CP	% of DM	19.0	17.0	16.0	15.0	14.0	13.0	16.0	
UIP	% of CP	40.0	35.0	32.0	30.0	25.0	25.0	30.0	
DIP ⁴	% of CP	60.0	65.0	68.0	70.0	75.0	75.0	70.0	
CP/ME	g CP/Mcal ME	65.0	66.0	65.0	63.0	59.0	55.0	61.0	
Minerals									
Ca	% of DM	.65	.55	.40	.30	.30	.30	.50	
P	% of DM	.45	.32	.30	.30	.25	.25	.35	
Mg ⁵	% of DM	.16	.16	.16	.16	.16	.16	.25	
K	% of DM	.65	.65	.65	.65	.65	.65	.65	
Salt		.30	.30	.30	.30	.30	.30	.30	
S	% of DM	.16	.16	.16	.16	.16	.16	.20	
Fe	ppm	50	50	50	50	50	50	50	
Co	ppm	.10	.10	.10	.10	.10	.10	.10	
Cu	ppm	10	10	10	10	10	10	10	
Mn	ppm	40	40	40	40	40	40	40	
Zn	ppm	40	40	40	40	40	40	40	
I	ppm	.25	.25	.25	.25	.25	.25	.25	
Se	ppm	.30	.30	.30	.30	.30	.30	.30	
Vitamins									
A	IU/lb of DM	1500	1000	1000	1000	1000	1000	2000	
D	IU/lb of DM	250	140	140	140	140	140	500	
E	IU/lb of DM	20	11	11	11	11	11	20	

¹ Assumes a calving age of 23 to 24 mo with heifers weighing 1400 lbs pre-calving.

² Prepuberty gains = 1.9 to 2.0 lbs/d; postpuberty gains = 1.7 to 1.8 lbs/d.

³ Energy levels may require reduction if ionophores are fed.

⁴ If DIP < 25.0% of CP, increase S to .25% of DM.

⁵ Additional Mg may be required when conditions favor tetany.

protein percent and amount of mammary tissue were not depressed by feeding a high energy, high gain (2.6 lbs/d) diet. Milk production results were, however, somewhat discouraging because the high energy, high protein diet resulted in significantly reducing calving age, but also reduced first lactation yield approximately 8.0 lbs/d. The reduction in milk yield cannot be well explained by mammary development reductions and likely is the result of body composition differences at calving. It should be noted that these trials contained rBST treatments, which help improve milk yields (59.3 lbs/d) of heifers fed the high energy, high protein diet. While not conclusive, these data (15, 16) suggest protein to energy ratios in heifer diets may be important.

Conclusions

Highlights from recent experiments regarding protein feeding of heifers are as follows:

- Calf starters require 19 to 20% CP on a DM basis (17 to 18% CP as fed).
- Calf starters should contain some high quality

bypass protein?

- Bypass protein recommendations in the 1989 NRC are too high.
- Replacement heifers do not consistently respond to supplemental bypass protein.
- Crude protein recommendations for heifers > 12 months of age in the 1989 NRC may be slightly conservative.
- When feeding high energy diets for accelerated growth, formulating diets on a CP to energy ratio may be of value.
- Optimizing rumen microbial protein synthesis is of importance in heifer diets (e.g., dietary carbohydrate inclusion).
- A data base on amino acid supplementation to replacement heifers and its economics is currently unavailable.

The author has attempted to summarize these new concepts in table 4.

(table 4. Ration specifications for Holstein replacement heifers)

References:

1. Abdelgadir, I. E. O., J. L. Morrill, and J. J. Higgins. 1996. Effect of roasted soybeans and corn on performance and ruminal and blood metabolites of dairy calves. *J. Dairy Sci.* 79:465-474.
2. Akayezu, J. M., J. G. Linn, D. E. Otterby, W. P. Hansen, and D. G. Johnson. 1994. Evaluation of calf starters containing different amounts of crude protein for growth of Holstein calves. *J. Dairy Sci.* 77:1882-1889.
3. Amos, H. E. 1986. Influence of dietary protein degradability and energy concentration on growth of heifers and steers and intraruminal protein metabolism. *J. Dairy Sci.* 69:2099-2110.
4. Bagg, J. G., D. G. Grieve, J. H. Burton, and J. B. Stone. 1985. Effect of protein on growth of Holstein heifer calves from 2 to 10 months. *J. Dairy Sci.* 68:2929-2939.
5. Bethard, G. L., R. E. James, and M. L. McGilliard. 1997. Effect of rumen-undegradable protein and energy on growth and feed efficiency of growing Holstein heifers. *J. Dairy Sci.* 80:2149-2155.
6. Casper, D. P., D. J. Schingoethe, M. J. Brouk, and H. A. Maiga. 1994. Nonstructural carbohydrate and undegradable protein sources in the diet: growth responses of dairy heifers. *J. Dairy Sci.* 77:2595-2604.
7. Heinrichs, J., and B. Lammers. 1998. Monitoring dairy heifer growth. *Ext. Circ. 5M498PS, Pennsylvania State Univ., University Park.*
8. Hoffman, P. C. 1997. Optimum body size of Holstein replacement heifers. *J. Anim. Sci.* 75:836-845.
9. Hoffman, P. C., N. M. Brehm, H. Chester-Jones, and M. Engstrom. 1998. Effect of dietary protein content on frame growth, blood metabolites, and nitrogen balance in Holstein heifers. *J. Dairy Sci.* 82(Suppl. 2): In press (Abstr.)
10. Hoffman, P. C., N. M. Brehm, S. G. Price, and A. Prill-Adams. 1996. Effect of accelerated post-pubertal growth and early calving on lactation performance of primiparous Holstein heifers. *J. Dairy Sci.* 79:2024-2031.
11. Luchini, N. D., S. F. Lane, and D. K. Combs.



-
1991. Evaluation of starter diet crude protein level and feeding regimen for calves weaned at 26 days of age. *J. Dairy Sci.* 74:3949-3955.
12. Mantysaari, P. E., C. J. Sniffen, T. V. Muscato, and M. L. Thonney. 1989. Performance of growing dairy heifers fed diets containing soybean meal or animal by-product meals. *J. Dairy Sci.* 72:2107-2114.
13. National Research Council. 1989. *Nutrient Requirements of Dairy Cattle*. 6th rev. ed. Natl. Acad. Sci., Washington, D. C.
14. Pirlo, G., M. Capelletti, and G. Marchetto. 1997. Effects of energy and protein allowances in the diets of prepubertal heifers on growth and milk production. *J. Dairy Sci.* 80:730-739.
15. Radcliff, R. P., M. J. VandeHaar, L. T. Chapin, T. E. Pilbeam, R. W. Ashley, S. M. Puffenbarger, E. P. Stanisiewski, D. K. Beede, and H. A. Tucker. 1998. Effects of diet and exogenous bST on reproductive performance on dairy heifers. *J. Dairy Sci.* 81(Suppl. 1):884.(Abstr.)
16. Radcliff, R. P., M. J. VandeHaar, A. L. Skidmore, L. T. Chapin, B. R. Radke, J. W. Lloyd, E. P. Stanisiewski, and H. A. Tucker. 1997. Effects of diet and bovine somatotropin on heifer growth and mammary development. *J. Dairy Sci.* 80:1996-2003.
17. Schwab, C. G. 1997. Unpublished data.
18. Schwartz, L. A., A. J. Heinrichs, G. A. Varga, and L. D. Muller. 1991. Effects of varying dietary undegradable protein on dry matter intake, growth, and carcass composition of Holstein calves. *J. Dairy Sci.* 74:3884-3890.
19. Steen, T. M., J. D. Quigley, III, R. N. Heitmann, and J. D. Gresham. 1992. Effects of lasalocid and undegradable protein on growth and body composition of Holstein heifers. *J. Dairy Sci.* 75:2517-2523.
20. Tomlinson, D. L., R. E. James, G. L. Bethard, and M. L. McGilliard. 1997. Influence of undegradability of protein in the diet on intake, daily gain, feed efficiency, and body composition of Holstein heifers. *J. Dairy Sci.* 80:943-948.
21. Van Amburgh, M. E., D. G. Fox, D. M. Galton, D. E. Bauman, and L. E. Chase. 1998. Evaluation of National Research Council and Cornell Net Carbohydrate and Protein Systems for predicting requirements of Holstein heifers. *J. Dairy Sci.* 81:509-526.
22. VandeHaar, M. J. 1998. Feeding heifers as a long-term investment. Page 1 in *Proc. Northwest Dairy Nutrition Shortcourse*. Jan. 30, 1998, Semiahmoo, WA, Washington State Univ.
23. Waldo, D. R., H. F. Tyrrell, A. V. Capuco, and C. E. Rexroad, Jr. 1997. Components of growth in Holstein heifers fed either alfalfa or corn silage diets to produce two daily gains. *J. Dairy Sci.* 80:1674-1684.

Notes

Notes



Western Dairy Management Conference • April 8-10, 1999 • Las Vegas, Nevada
